

Flash Sfil

Catalog No.: 2527-100 Size: 100 rxns

5'...G G C C N N N N N G G C C...3'
 3'...C C G G N N N N N C C G G...5'



Storage Condition

-20°C

Components

Components	Amount
Flash Sfil (20 U/μl)	100 μl (2000 U)
10× Flash Buffer	1 ml
10× Flash Color Buffer	1 ml

Description

Flash enzymes are a series of engineered restriction enzymes that are capable of fast DNA digestion. All Flash enzymes show superior activity in the universal Flash and Flash Color Buffer, and are able to digest DNA in 5~15 minutes. This enables any combination of restriction enzymes to work simultaneously in one reaction tube and eliminates the need for sequential digestions. Flash enzymes have passed multiple strict quality controls, and can be used to digest plasmid, genomic and viral DNA as well as PCR products.

Flash Color Buffer includes a density reagent along with red and yellow tracking dyes that allow for direct loading of the reaction mixtures on a gel. The red dye of the Flash Color Buffer migrates with 2.5 kb double-strand DNA fragments in a 1% agarose gel, and the yellow dye migrates with 10 bp double-strand DNA fragments in a 1% agarose gel.

Recommended Reaction Conditions

1× Flash Buffer;
 Incubate at 50°C ;

Refer to "Protocol for Fast DNA Digestion" for reaction setup.

Heat Inactivation

This enzyme can not be heat inactivated. Please purify the digested product by phenol/chloroform treatment or column-based purification kit.

Quality Control

Functional Test

A 20 μl reaction in Flash Buffer containing 1 μg of pEPE DNA and 1 μl of Flash Sfil incubated for 15 minutes at 50 °C results in complete digestion as determined by agarose gel electrophoresis.

Prolonged Incubation / Star Activity Assay

A 20 μl reaction in Flash Buffer containing 1 μg of pEPE DNA and 1 μl of Flash Sfil incubated for 3 hours at 50°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. Longer incubation may result in star activity.







Ligation and Recutting

After 10-fold over-digestion with Flash Sfil at 50°C , >95% of the DNA fragments can be ligated with T4 DNA Ligase at 22 °C . Of these ligated fragments, >95% can be recut with Flash Sfil as determined by agarose gel electrophoresis.

Non-specific Endonuclease Activity

A 20 μl reaction in Flash Buffer containing 1 μg of supercoiled plasmid and 1 μl of Flash Sfil incubated for 4 hours at 50°C results in <10% conversion to the nicked or linearized form as determined by agarose gel electrophoresis.

Icon Descriptions

-  This enzyme will digest unit substrate in 5~15 minutes under recommended reaction conditions.
-  The enzyme's optimum reaction temperature is 50°C .
-  Cleavage with this restriction enzyme may be blocked or impaired when the substrate DNA is methylated by the CpG methylase.
-  Cleavage with this restriction enzyme may be blocked or impaired when the substrate DNA is methylated by the Dcm methylase.
-  The enzyme can not be thermal inactivated.
-  3 hours incubation do not show star activity, but longer incubation may result in star activity.

Method of application

1. Protocol for Fast DNA Digestion

① Combine the following reaction components on ice in the order indicated:

	Plasmid DNA	PCR product	Genomic DNA
ddH ₂ O	15 µl	16 µl	30 µl
10× Flash Buffer or 10× Flash Color Buffer	2 µl	3 µl ^a	5 µl
DNA	2 µl (up to 1 µg)	10 µl (~0.2 µg)	10 µl (5 µg)
Flash Sfil	1 µl	1 µl	5 µl
Total	20 µl	30 µl	50 µl

a. For purified PCR products. If the PCR products are not purified, amount of 10× Flash Buffer should be reduced to 2 µl due to the remaining metal ions in the unpurified PCR products. We recommend to purify PCR products before digestion if it will be used for cloning, because the exonuclease activity of some DNA polymerases may alter the end of cleaved DNA.

- ② Mix gently and spin down;
- ③ Incubate at 50°C for 15 minutes (plasmid DNA) or for 15~30 minutes (PCR product) or for 30~60 minutes (genomic DNA);
- ④ Optional: Phenol/chloroform treatment or column based purification.
- ⑤ If the Flash Color Buffer was used in the reaction, load an aliquot of the reaction mixture directly on a gel.

2. Double and Multiple Digestion of DNA

- ① Use 1 µl of each enzyme and scale up the reaction conditions appropriately;
- ② The combined volume of the enzymes in the reaction mixture should not exceed 1/10 of the total reaction volume;
- ③ If the enzymes require different reaction temperatures, start with the enzyme that requires a lower temperature, then add the second enzyme and incubate at the higher temperature.

3. Scaling up Plasmid DNA Digestion Reaction

DNA	1 µg	2 µg	3 µg	4 µg	5 µg
Flash Sfil	1 µl	2 µl	3 µl	4 µl	5 µl
10× Flash Buffer or 10× Flash Color Buffer	2 µl	2 µl	3 µl	4 µl	5 µl
Total	20 µl	20 µl	30 µl	40 µl	50 µl

✦ Note: Increase the incubation time if the total reaction volume exceeds 20 µl.

Number of Recognition Sites in DNA

λDNA	ΦX174	pBR322	pUC57	pUC18/19	SV40	M13mp18/19	Adeno2
0	0	0	0	0	1	0	3

Methylation Effects on Digestion

Dam	Dcm	CpG	EcoKI	EcoBI
No effect	Impaired	Impaired	No effect	No effect

Activity in Different Buffers*

	Flash Buffer	Thermo Scientific FastDigest Buffer	NEB rCutSmart™ Buffer	Takara QuickCut™ Buffer
Activity	100%	100%	100%	100%

*The activity data come from the functional test described above.

Activity of DNA Modifying Enzymes in Flash and Flash Color Buffers

2601-01K Alkaline Phosphatase (Flash)	100%
2600-01K T4 DNA Ligase (Flash)*	100%

*ATP is required for T4 DNA Ligase activity.